

## 3.A3.B Worksheet B – Initial Amplification of Nucleic Acid by PCR for the Confirmation of *Renibacterium salmoninarum*

Case Number \_\_\_\_\_

Date \_\_\_\_\_

PCR Reagent	Lot#	Final Concentration	Stock Concentration**	Volume per Reaction (µL) (to total 50 µL)	Volume for ___ samples
d-H <sub>2</sub> O*		Add to total 40 µL		30.1	
10XBuffer		1X	10X	5.0	
MgCL <sub>2</sub>		1.5 mM	50 mM	1.5	
dNTP's		0.2 mM	10 mM	1	
(+)Primer		20 pMole	20 pMole/µL	1	
(-)Primer		20 pMole	20 pMole/µL	1	
TAQ		2 units/Rx	5U/µL	0.4	
DNA <sup>‡</sup>		-	-	10 µL	-

\*Add water to Master Mix first, TAQ last.

\*\*Change “Stock Concentration” parameters as necessary. Different reagent sources supply varying stock concentrations.

‡Do not add DNA template until Master Mix reaction tubes have been removed from the reagent mixing (MM) area.

### Primer Sets for *R. salmoninarum* 1<sup>st</sup> Round Amplification

<b>P3 (round 1 forward)</b>	5'-A GCT TCG CAA GGT GAA GGG-3'
<b>M21 (round 1 reverse)</b>	5'-GC AAC AGG TTT ATT TGC CGG G-3'