

- hole ecosystems and microcosms. *Ecology* 72:1529-1546.
- Wallace, J. B., and R. W. Merritt. 1980. Filter-feeding ecology of aquatic insects. *Annual Review of Entomology* 25:103-132.
- Wharton, G. W. 1985. Water balance of insects. p. 565-601. In G. A. Kerkut, and L. I. Gilbert (eds.) *Comparative Insect Physiology, Biochemistry, and Pharmacology*, Vol. 3. 1st ed. Pergamon Press, Oxford.
- Wigglesworth, V. B. 1938. Regulation of osmotic concentration and chloride in blood of mosquito larvae. *Journal of Experimental Biology* 15:235-247.

Mangum, F.A. and J.L. Madrigal
Rotenone Effects on Aquatic Invertebrates
of The Strawberry, Utah: A Five-year Summary

USDA Forest Service National Aquatic Ecosystem Analysis Lab
Uinta National Forest 88 West 100 North
Provo, UT 84601 USA

and

J. L. Madrigal
222 TMCB, Department of Statistics
Brigham Young University
Provo, UT 84601 USA

ABSTRACT

Before treatment with a 3 mg/l Noxfish (0.15 mg/l active ingredient; rotenone) for 48 hours, benthic invertebrate communities were quantitatively sampled with a modified Surber net. Then spring, summer, and fall post-rotenone samples were taken monthly at each of four Strawberry River stations for five years. Statistical analyses of the data indicated that the application of rotenone had a significant effect on the following species density: *Cinygmula* sp., *Pteronarcella badia*, *Hesperoperla pacifica*, *Hydropsyche* sp., and *Brachycentrus americanus*. Thirty-three percent of the benthic invertebrate taxa at the four stations showed resistance to rotenone. Up to 100% of Ephemeroptera, Plecoptera and Trichoptera species were missing after the second rotenone application. Forty-six percent of the taxa recovered within one year, but 21% of the taxa were still missing after five years. Of the 19 taxa still missing, 47% were Trichoptera, 21% were Ephemeroptera, 16% were Plecoptera, 11% were Coleoptera, and 5% were Megaloptera.

INTRODUCTION

It is common in management of certain fisheries programs to eradicate all nonnative or exotic species from a freshwater habitat so that desired native species can be given a noncompetitive fresh start in an aquatic ecosystem. This method has generally been used in recovery programs for threatened or endangered fish species. One of the most popular ways to remove fish is by treating aquatic ecosystems with rotenone. The primary action of rotenone is to block important biochemical pathways of cell metabolism (Lindahl and Oberg 1961, Oberg 1962). Rotenone inhibits the respiration of mitochondria by blocking the reduced nicotinamide adenine dinucleotide (NADH)-dehydrogenase segment of the respiratory chain in fish and aquatic insects (Fukami, et al. 1969). Because of this nonspecific poisoning, many non target species such as certain aquatic macroinvertebrates are also eliminated from the ecosystem.

Fish toxicants have generally been applied without specific knowledge concerning possible adverse effects on aquatic macroinvertebrates living in the ecosystems treated. Gilderhaus, et al. (1988) used high performance liquid chromatography to analyze rotenone concentrations in water, bottom sediments, invertebrates, and fish

none treatment. Decomposition of rotenone in water followed a first-order decay curve; half life was 10.3 days in cold water (0-10°C) and 0.94 days in warm water (23-27°C). The effects of rotenone on aquatic invertebrates have been recorded by Binns (1967), Cook and Moore (1969), and Engstrom-Heg, et al. (1978). Each of these investigators observed that stream bottom insects varied greatly in their tolerances to rotenone.

We compared pre-rotenone benthic invertebrate communities and taxa to post-rotenone community composition for a five-year period following rotenone treatment of tributaries to Strawberry Reservoir in Utah. This study included more specific taxonomy than found in previous studies. Many taxa were classified to the genus and species levels to determine more specifically the effects of rotenone on species of aquatic insects and other invertebrates in treated streams. This is also the first time recovery or lack of recovery of benthic community members has been followed for a five-year period. The Strawberry River rotenone project, which was the largest ever attempted in the U.S.A., included a 5,666 HA reservoir and 274 km. of perennial tributary streams. The rotenone treatments were applied to remove Utah Chubs (*Gila atraria*) and Utah Suckers (*Catostomus ardens*) from the Strawberry Reservoir and drainage. The target fish species were competing with the trout in the reservoir and tributaries. In the Strawberry ecosystem, three salmonid species were introduced into post-treatment waters. These included sterile *Oncorhynchus mykiss* and Bear Lake *Salmo clarki* to help control rough fish, and *Oncorhynchus nerka*, a noncompetitive plankton feeder.

Aquatic ecosystem quality varied between stations included in this study. The pre-rotenone aquatic macroinvertebrate communities at the lower Strawberry River stations were dominated by tolerant species before the rotenone project began. The upper two stations had more sensitive species in the pre-rotenone benthic communities (Mangum 1995).

MATERIALS AND METHODS

Rotenone Application

Rotenone was applied in 1990 to the entire Strawberry River watershed (Fig. 1). The rotenone used was Noxfish, which had a 5% by volume active ingredient. Drip barrels were set up to apply 3 mg/l Noxfish on the streams. The goal was to maintain a 0.15 mg/l active ingredient concentration in the stream channels for 48 hours. To provide unrestricted flows, beaver dams were breached the day prior to treatment. Ground crews used backpack sprayers to apply liquid rotenone to backwater and side channel areas not reached by flows carrying the rotenone from drip barrels. A powdered rotenone/sandmix mixture was applied to seeps, springs, and weedy areas. All streams, springs, and seeps were treated to the tops of the headwaters in the entire Strawberry Reservoir drainage. Rotenone was applied twice. The first application was in early to mid-August; the second one was applied from September 25 through October 16.

Aquatic Macroinvertebrate Sampling

Three quantitative macroinvertebrate samples were taken monthly with a one meter long, 280 micron mesh, 45.7 cm upper frame Winget-modified Surber net (Winget 1979). This was done from spring (May to

June) through fall (September to October) at each station each year.

At Strawberry River stations (Fig. 1), pre-rotenone communities were a composite of taxa present in June and early August 1990. Samples were taken about one week prior to treatment, and again seven to ten days following each of the two rotenone applications. Since ethanol (80%) was used to preserve the organisms, if samples were taken too soon after the rotenone application, it was difficult to determine if the organisms died from rotenone or ethanol. Thus, macroinvertebrates allowed to decay for seven to ten days following rotenone treatment were easily distinguished and were not counted as survivors.

Unprocessed portions of subsamples were scanned for taxa that may not have been included in sorted subsamples. Aquatic macroinvertebrate nymphs and larvae were classified to the species or genus level when possible so that tolerance levels and recovery could be followed more specifically.

Statistical Analysis

The data were analyzed using a generalized linear model approach. This technique is recommended when the response variable is not normal, and/or when the variance is not constant (McCullagh and Nelder, 1989). Construction of a generalized linear model requires choosing an appropriate link function and response probability distribution. The data for this project were modeled using a binomial error distribution with a probit link function. This modeling process was used for each taxon. It is important to mention that station and date were treated as explanatory variables. In all cases, the research hypothesis was:

H₀: Rotenone does not have any effect on a species density

H_a: Rotenone has a significant effect on a species density

The Procedure GENMOD from the statistical software SAS® was utilized to test the hypothesis. The hypothesis was rejected when the p-value was smaller than 0.05.




Figure 1. Location of sampling sites in the Strawberry River system, Utah.

RESULTS

The statistical analysis was limited to the six more important species (*Drunella doddsi*, *Cinygmula* sp., *Pteronarcella badia*, *Hesperoperla pacifica*, *Hydropsyche* sp., and *Brachycentrus americanus*). The results showed significant differences by station and date in the densities (due to rotenone application, p -value=0.0001) of all of the above species except *Drunella doddsi*. A similar behavior was observed through time. In other words, their population densities decreased considerably due to rotenone.

The first rotenone application eliminated many of the Ephemeroptera, Plecoptera, and Trichoptera (EPT) species from the community. Populations of Diptera and Oligochaeta taxa were initially reduced, but post-rotenone conditions seemed to favor these taxa. Within one to two months after treatment the communities were dominated by simuliids, chironomids, Oligochaeta, and Tipula.

At the four Strawberry River Stations, up to 59% of the taxa were removed from the communities by the first rotenone application. The second rotenone application reduced the pre-rotenone communities by up to 73%.

After the first rotenone application, the benthic communities had 45-82% fewer Ephemeroptera, 50-69% fewer Plecoptera, and 30-75% fewer Trichoptera. Although numbers of most other taxa were obviously reduced, they were not eliminated from the communities. After the second

Table 1. Post-rotenone recovery of Strawberry River taxa - Station 1

Recovery Time	Taxa Recovered
0 months (Not Missing)	<i>Tricorythodes minutus</i> , <i>Paraleptophlebia</i> sp., <i>Hydropsyche</i> sp., <i>Cheumatopsyche</i> sp., <i>Helicopsyche borealis</i> , <i>Oecetis</i> sp., <i>Optioservus</i> sp., <i>Zaitzevia</i> sp., <i>Tipula</i> sp., Chironomidae, <i>Chelifera</i> sp., <i>Bezzia</i> sp., <i>Physsa</i> sp., Oligochaeta, Nematoda, <i>Hydracarina</i> sp.
2 months	<i>Baetis</i> sp., Simuliidae
9-12 months	<i>Heptagenia</i> sp., <i>Drunella grandis</i> , <i>Ephemerella inermis</i> , <i>Sweltsa</i> sp., <i>Isoperla fulva</i> , <i>Pteronarcella badia</i> , <i>Arctopsyche</i> sp., <i>Dicosmoecus</i> sp., <i>Limnephilus</i> sp., <i>Brachycentrus occidentalis</i> , <i>Hydroptila</i> sp., <i>Protophila</i> sp., <i>Agabus</i> sp., <i>Hexatoma</i> sp., <i>Euparyphus</i> sp., Planorbidae, <i>Lymnaea</i> sp., Pelecyopoda, <i>Gammarus</i> sp.
21 months	<i>Leucotrichia</i> sp.
34 months	<i>Helobdella</i> sp.
58 months	<i>Ochrotrichia</i> sp.
Still missing after 5 years	<i>Cinygmula</i> sp., <i>Atenuatella margarita</i> , <i>Caudatella hystrix</i> , <i>Rhyacophila acropedes</i> , <i>Agapetus</i> sp., <i>Hydaticus</i> sp.

rotenone application, Ephemeroptera diversity was reduced by 67-100%, Plecoptera by 67-100%; and Trichoptera by 61-100%. The percent of taxa, by order, missing after one year were: 20-45% of the Ephemeroptera, 20-100% of the Plecoptera and 20-100% of the Trichoptera (Tables 1-4).
Tolerances and Recovery of Taxa

In this study, a taxon was considered missing if it was no longer found in the community after the first or second application of rotenone. As soon as a taxon reappeared in the sampled community at a given station, it was considered to have recovered.

At Station 1, of the 46 taxa in the pre-rotenone community, 15 taxa were resistant to rotenone. Of the 31 missing after rotenone application, two recovered within two months and twenty others recovered after 12 months. One taxon recovered within 21 months and another recovered in the third year. After five years, six taxa found in the pre-rotenone community were still missing (Table 1).

Brown and Ball (1943) state that leeches were seriously affected by a 0.5 mg/l concentration of 5% *Derris* powder. Leeches in this study recovered after 34 months at Strawberry River Station 1.

At Station 2, of the 56 taxa in the pre-rotenone community, five were resistant to rotenone. Three recovered within two months, twenty more recovered within 12 months, two more by 14 months, six took 19-24 months to recover, and another taxon recovered within 35 months. After five years, eight taxa were still missing (Table 2).

Table 2. Post-rotenone recovery of Strawberry River taxa - Station 2

Recovery Time	Taxa Recovered
0 months (Not Missing)	<i>Arctopsyche</i> sp., <i>Tipula</i> sp., <i>Hexatoma</i> sp., <i>Euparyphus</i> sp., <i>Lymnaea</i> sp.
2 months	Simuliidae, <i>Hemerodromia</i> sp., <i>Chelifera</i> sp.
8-12 months	<i>Rhithrogena</i> sp., <i>Heptagenia</i> sp., <i>Drunella Grandis</i> , <i>Ephemerella inermis</i> , <i>Paraleptophlebia</i> sp., <i>Baetis</i> sp., <i>Sweltsa</i> sp., <i>Pteronarcella badia</i> , <i>Isoperla</i> sp., <i>Hydropsyche</i> sp., <i>Hydroptila</i> sp., <i>Rhyacophila vagrita</i> , <i>Wormaldia</i> sp., <i>Leucotrichia</i> sp., <i>Hesperophylax</i> sp., <i>Agabus</i> sp., Carabidae, <i>Antocha monticola</i> , Planorbidae, Pelecyopoda
13-14 months	<i>Cinygmula</i> sp., <i>Hydaticus</i> sp.
19-24 months	<i>Tricorythodes minutus</i> , <i>Hesperoperla pacifica</i> , <i>Brachycentrus americanus</i> , <i>Helicopsyche borealis</i> , <i>Dicosmoecus</i> sp., Ostracoda
35 months	<i>Serratella tibialis</i>
60-61 months	Capniidae, <i>Pericoma</i> sp.
Still missing after 5 years	<i>Caudatella hystrix</i> , <i>Isogenoides</i> sp., <i>Brachycentrus occidentalis</i> , <i>Micrasema</i> sp., <i>Ochrotrichia</i> sp., <i>Glossosoma</i> sp., <i>Oecetis</i> sp., <i>Stalis</i> sp.

At Station 3, of the 59 taxa in the pre-rotenone community: 16 appeared to be resistant to rotenone. Of the 43 taxa missing after rotenone treatments, twenty recovered within 12 months. Thirteen others recovered within the second year, two more recovered in the third year, and one recovered in the fourth year (Table 3). After five years, there were five taxa still missing (Table 3).

Of the 55 taxa in the pre-rotenone community at Station 4, 16 appeared to be resistant to the rotenone concentration used at this station. Of the 38 taxa missing after rotenone applications, twenty nine recovered within 12 months and one after 14 months. After one year, 16% of the pre-rotenone community was still missing. Four taxa recovered within the second year, and another taxon recovered within 25 months. After five years, four species were still missing (Table 4).

Table 3. Post-rotenone recovery of Strawberry River taxa - Station 3

Recovery Time	Taxa Recovered
0 months (Not Missing)	<i>Paraleptophlebia</i> sp., <i>Hydropsyche</i> sp., <i>Optioservus</i> sp., <i>Zaitzevia</i> sp., <i>Hexatoma</i> sp., <i>Tipula</i> sp., Simuliidae, Chironomidae, <i>Chelifera</i> sp., <i>Bezzia</i> sp., <i>Pericoma</i> sp., <i>Euparyphus</i> sp., Ostracoda, <i>Oligochaeta</i> , <i>Hydracarina</i> sp., Nematoda
8-12 months	<i>Heptagenia</i> sp., <i>Drunella doddsi</i> , <i>Drunella grandis</i> , <i>Baetis</i> sp., <i>Cultus</i> sp., <i>Amphinemura</i> sp., <i>Podmosta besameta</i> , Capniidae, <i>Eucapnopsis</i> sp., <i>Isoperla fulva</i> , <i>Hesperoperla pacifica</i> , <i>Sweltsa coloradensis</i> , <i>Arctopsyche grandis</i> , <i>Dicosmoecus</i> sp., <i>Hydroptila</i> sp., <i>Rhyacophila hyalinata</i> , <i>Rhyacophila vagrita</i> , <i>Agabus</i> sp., <i>Atrichopogon</i> sp., <i>Linnophora</i> sp.
20-24 months	<i>Epeorus longimanus</i> , <i>Rhithrogena hageni</i> , <i>Serratella tibialis</i> , <i>Zapada haysi</i> , <i>Kogotus modestus</i> , <i>Plumiperla diversa</i> , <i>Hesperophylax</i> sp., <i>Helicopsyche borealis</i> , <i>Rhyacophila acropedes</i> , <i>Lepidostoma</i> sp., <i>Antocha monticola</i> , <i>Dicranota</i> sp., <i>Sialis</i> sp.
32-36 months	<i>Drunella coloradensis</i> , <i>Suwallia pallidula</i> , <i>Dixa</i> sp.
47 months	<i>Brachycentrus americanus</i>
Still missing after 5 years	<i>Caudatella hystrix</i> , <i>Podmosta delicatula</i> , <i>Isogenoides</i> sp., <i>Oligophlebodes</i> sp., Carabidae

DISCUSSION

Cushing and Olive (1957) reported that *Oligochaeta* increased after rotenone treatment. In the Strawberry River study, populations of chironomids and tubificids were adversely affected but recovered to twice their original numbers within one month. Post-rotenone increases in benthic invertebrate populations may be due in part to elimination of fish

and other predators (Tuunainen 1970). Hubbs (1963), observed that application of rotenone on the Concho River in Texas drastically altered the ecological interactions and had selective effects, particularly on game fish food organisms. Little (1966) reported that 34% to 100% of bottom-dwelling organisms were removed by rotenone treatment. He found that all organisms except Plecoptera and *Isopoda* recovered up to their original abundance within one year.

Binns (1967) found that after application of rotenone, taxonomic groups recovering in twelve months included Tipulidae, Tricorythodidae, Heptageniidae, Perlidae, and Lepidostomatidae. Brachycentridae and Hydropsychidae took 20 months, Oligochaeta took 15 months, and Empididae took 24 months. Baetidae recovered in three months. Recovery was probably assisted by organisms drifting from upstream, untreated reaches.

Some of the possible mechanisms for recovery or survival from effects of rotenone treatment could include survival of adults, nymphs, or larvae of species physiologically or structurally resistant to the adverse effects of rotenone. An example of a resistant nymph is *Paraleptophlebia* sp. At most stations on the Strawberry River, its numbers remained consistently high before, during, and after the rotenone application. This genus was observed to be tolerant to rotenone by Engstrom-Heg et al. (1978); they also found the caddisfly *Hydropsyche* sp. Was tolerant to

Table 4. Post-rotenone recovery of Strawberry River taxa - Station 4

Recovery Time	Taxa Recovered
0 months (Not Missing)	<i>Ephemerella inermis</i> , <i>Paraleptophlebia</i> sp., <i>Baetis</i> sp., <i>Rhyacophila acropedes</i> , <i>Rhyacophila vagrita</i> , <i>Zaitzevia</i> sp., <i>Optioservus</i> sp., Chironomidae, <i>Hemerodromia</i> sp., <i>Bezzia</i> sp., <i>Pericoma</i> sp., <i>Euparyphus</i> sp., Ostracoda, <i>Hydracarina</i> sp., <i>Oligochaeta</i> , Nematoda
8-12 months	<i>Epeorus</i> sp., <i>Cinygmula</i> sp., <i>Drunella coloradensis</i> , <i>Drunella doddsi</i> , <i>Tricorythodes minutus</i> , Chloroperlidae, <i>Sweltsa</i> sp., <i>Megarcys signata</i> , <i>Isoperla</i> sp., <i>Cultus</i> sp., <i>Isogenoides</i> , <i>Zapada haysi</i> , <i>Zapada cinctipes</i> , <i>Hesperoperla pacifica</i> , Leuctridae, <i>Arctopsyche grandis</i> , <i>Lepidostoma</i> sp., <i>Rhyacophila hyalinata</i> , <i>Rhyacophila vepulsa</i> , <i>Dicosmoecus</i> sp., <i>Neothremma</i> sp., <i>Hesperophylax</i> sp., <i>Hexatoma</i> sp., <i>Dicranota</i> sp., Simuliidae, <i>Chelifera</i> sp., <i>Atrichopogon</i> sp., <i>Dixa</i> sp., <i>Planaria</i> sp.
14 months	<i>Tipula</i> sp.
20-24 months	<i>Rhithrogena hageni</i> , <i>Serratella tibialis</i> , <i>Skwala americana</i> , <i>Pelecypoda</i>
Still missing after 5 years	<i>Drunella grandis</i> , <i>Suwallia</i> sp., <i>Podmosta delicatula</i> , <i>Parapsyche elsis</i>

rotenone. We observed the same for *Hydropsyche* sp. in the Strawberry River and suggest that instream macrophytes could provide the needed oxygen or advantage for survival of this and other benthic species.

Some of the survivors in the Strawberry River may have been associated with instream or side springs which could locally dilute the rotenone. Also, there may have been incomplete mixing of the rotenone in some stream areas such as under banks or under rocks.

Within a week following the rotenone treatments there were few surviving benthic invertebrates observed in the samples, but within one to two months, populations of chironomids and simuliids were high. When compared with pre-rotenone sampled populations, simuliids at Station 1 had increased 240% by October 1990 and 1,108% by June 1991. Chironomids at Station 2 increased 232% by August 1991.

Many of the aquatic macroinvertebrates in the Strawberry tributaries that reappeared within a year following rotenone application may have been from eggs deposited in the stream before the project. Eggs are generally more resistant to adverse conditions than are nymphs or larvae, which are more dependent upon a sustained source and use of oxygen. Even fish eggs are more resistant to toxicants than young or mature fish (Lennon et al. 1970). It appears that some of the recovering species may have been reintroduced into the aquatic ecosystem through mechanisms such as wind current carrying aerial adults.

Most of the species sensitive to rotenone were in the orders Ephemeroptera, Plecoptera, and Trichoptera. (EPT). This was shown by the high mortality rate (up to 100%) for these taxa after the rotenone applications at each station. Twenty-two to 53% of these taxa recovered within one year, but 7 to 14% of the species were still missing after five years. Most of the aquatic macroinvertebrate species that are sensitive to different types of environmental perturbations are in the EPT orders. EPT species were the most sensitive to rotenone, but some recovered more quickly than expected. Recovery of some aquatic macroinvertebrate species could be delayed by ecosystem stress from past or current cattle or sheep grazing activities.

Because of the differences in habitat quality at the stations sampled, the benthic invertebrate community composition was different at each station monitored; however, some of the same species were still missing from the sampled communities after five years. *Caudatella hystrix* was still missing at Stations 1, 2, and 3, *Oecetis* sp and *Sialis* sp. at Station 2 on the Strawberry River, and *Podmosta delicatula* at Stations 3 and 4. Binns (1967) observed that none of the taxonomic groups were missing at the end of a two-year Green River study. However, more specific taxonomy used in the present study revealed that some taxa were still missing after five years. Long-term effects of rotenone were most evident for Trichoptera. Of the 19 taxa still missing at the four stations, 21% were Ephemeroptera, 16% were Plecoptera, 47% were Trichoptera, 11% were Coleoptera, and 5% were megalopterans.

Aquatic invertebrate taxa that were tolerant to the rotenone concentration used are identified as not-missing or briefly missing in the station taxa lists. Nine percent to 33% of the benthic invertebrates at the stations sampled were resistant to rotenone. Of the 27 resistant taxa, 11% were Ephemeroptera, none were Plecoptera, 26% were Trichoptera, 8%

were Coleoptera, 33% were Diptera and 22% were other invertebrates. Members of the family Simuliidae (blackflies) compete best and thrive where there is organic nutrient loading (Mangum 1995). The riffle beetles (Elmidae) and a mayfly, *Tricorythodes* sp., are tolerant to fine sediments (Mangum 1995). Many species in the dipteran family Chironomidae, the mayfly family Baetidae and aquatic worms (Oligochaeta) compete best and can become dominant in communities where there is an abundance of sediment and organic enrichment. Other tolerant taxa include the water mites (*Hydracarina* sp.).

Binns (1967) observed that after rotenone treatment the dominant benthos at the various stations were water beetles, midges, crane fly larvae, Perlodidae, Rhagionidae, Simuliidae, and *Tricorythodes* sp.. We found immediate changes in benthic invertebrate community composition and dominance, and there were changes in sets of dominant taxa during the five-year post-rotenone recovery period. Some of the taxa such as *Tipula* sp. and *Paraleptophlebia* sp. which were relatively obscure in diverse pre-rotenone communities became dominant survivors in post-rotenone communities. Numbers of benthic invertebrate organisms found in pre-rotenone samples were reestablished in one to 36 months. Binns (1967) similarly observed that although numbers of organisms were drastically reduced after rotenone applications, numbers soon became high again but dominance in the communities changed.

Not only are aquatic invertebrate species affected but the nutrient cycle and food chain components can be changed by rotenone. Some species not found in pre-rotenone communities were collected in post-rotenone samples. It will be of interest to see if they permanently fill the niches of missing species. We found that rotenone can cause long-term changes in aquatic macroinvertebrate community stability and composition.

We suggest that pre and post-treatment monitoring should be a part of any project using rotenone as a management tool. Rotenone concentrations used should be just enough to remove the target fish species, which could limit the number of non-target aquatic invertebrates eliminated by the treatment. To prevent loss of sensitive aquatic invertebrates, representative species of communities from stream reaches to be treated should be collected and placed in a suitable habitat and then returned to the stream reaches after rotenone toxicity subsides.

ACKNOWLEDGEMENTS

We wish to extend thanks to the following: Peter Karp, USDA Forest Supervisor; Barbara Franano and Paul Skabelund for forest program planning and assistance in collection of data; Joel Montgomery for field collections; Richard Baumann for stonefly identifications; Roger Wilson and Bob Spateholts, Utah Division of Wildlife Resources Biologists, for providing information on rotenone technology used; Brian Johnston, USFS Lab Typist; Nicole Cox, USFS Aquatic Lab Senior Technician, for insect identifications and manuscript review; Don Duff, USFS/TU Aquatic Ecologist for manuscript review.

LITERATURE CITED

- Binns, N. A. 1967. Effects of rotenone treatment on the fauna of the Green River, Wyoming. Wyoming Game and Fish Commission--Fisheries Research Bulletin 1:1-108.
- Cook, S. F., Jr., and R. L. Moore. 1969. The effects of rotenone treatment on the insect fauna of a California stream. American Fisheries Society, Trans. 98(3): 539-544.
- Cushing, C. E., Jr., and J. R. Olive. 1957. Effects of toxaphene and rotenone upon the macroscopic bottom fauna of two northern Colorado reservoirs. American Fisheries Society, Trans. 86:294-301.
- Engstrom-Heg, R., R. T. Colesante, and E. Silco. 1978. Rotenone tolerances of stream-bottom insects. New York Fish and Game Journal 25 (1):31-41.
- Fukami, J.-I., T. Shishido, K. Fukunaga, and J. E. Casida. 1969. Oxidative metabolism of rotenone in mammals, fish, and insects and its relation to selective toxicity. J. Agr. Food Chem.--US Public Health Service grant ESGM 00049 and AEC Contract AT (11-1) 34, Project Agree No. 113, 1217-1226 p.
- Gilderhaus, P. A., V. K. Dawson, and J. L. Allen. 1988. Deposition and persistence of rotenone in shallow ponds during cold and warm seasons. USDI Fish and Wildlife Service--National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161, 7 pp.
- Hubbs, C. 1963. An evaluation of the use of rotenone as a means of "improving" sports fishing in the Concho River, Texas. Copeia, 1:99-203.
- Lennon, R. E., J. B. Hunn, R. A. Schnick, and R. M. Burress. 1970. Reclamation of ponds, lakes, and streams with fish toxicants: A review. USDI, Fish and Wildlife Service-FAO Fisheries Technical paper 100. FIRI/T100. Inland Resources Management. 99 pp.
- Lindahl, P. E., and K. E. Oberg. 1961. The effect of rotenone on respiration and its point of attack. Experimental Cell Research 23:228-237.
- Lindgren, P. E. 1960. About the effect of rotenone upon benthonic animals in lakes. Fisheries Board of Sweden. Report of the Institute of Freshwater Research, Drottningholm 41:172-184.
- Little, J. D. 1960. About the effect of rotenone upon benthic animals in lakes. Institute of Freshwater Research, Drottningholm. Report No. 41:172-184.
- Mangum, F. A. 1995. Aquatic Ecosystem Inventory-Macroinvertebrate Analysis-Uinta National Forest Strawberry Tributaries. 153 pp.
- McCullagh, P. And Nelder, J.A. (1989) *Generalized Linear Models*, London: Chapman and Hall
- Oberg, K. E. 1962. The site of action of rotenone in the respiratory chain. Experimental Cell Research 24:163-164.
- Pennak, R. W. 1953. Freshwater invertebrates of the United States. New York: Ronald Press.
- Tuunainen, P. 1970. Relations between the benthic fauna and two species of trout in some small Finnish lakes treated with rotenone. Annales Zoologici Fennici 7:67-120.
- Winget, R.N. and F.A. Mangum. 1979. Biotic Condition Index: Integrated Biological, Physical, and Chemical Stream parameters for Management. USDA Forest Service, Intermountain Region.
- Wollitz, R. E. 1962. Effects of certain commercial fish toxicants on the limnology of three cold-water ponds, Montana. Proceedings of the Montana Academy of Sciences 22:54-81.